

Table 7. Estimated numbers of deaths of persons with AIDS, by year of death and selected characteristics, 2000–2004—United States

	Year of death					Cumulative through 2004 ^a
	2000	2001	2002	2003	2004	
Age at death (yrs)						
<13	52	46	32	28	18	5,094
13–14	8	4	11	7	16	266
15–19	41	44	39	43	32	1,055
20–24	167	215	167	175	184	8,808
25–29	710	635	595	569	505	44,516
30–34	1,993	1,744	1,555	1,373	1,157	96,357
35–39	3,346	3,292	3,108	2,969	2,404	116,206
40–44	3,523	3,835	3,726	3,800	3,378	100,633
45–49	3,081	3,121	3,364	3,432	3,016	67,842
50–54	1,966	2,152	2,396	2,524	2,314	39,936
55–59	1,007	1,141	1,228	1,403	1,343	22,452
60–64	593	655	621	726	701	12,946
≥65	652	728	702	801	730	13,004
Race/ethnicity						
White, not Hispanic	5,325	5,194	5,210	5,091	4,316	229,220
Black, not Hispanic	8,605	9,011	8,974	8,950	7,978	201,045
Hispanic	3,025	3,195	3,117	3,537	3,228	93,163
Asian/Pacific Islander	95	100	91	81	82	3,272
American Indian/Alaska Native	66	81	84	73	91	1,578
Transmission category						
Male adult or adolescent						
Male-to-male sexual contact	5,955	6,068	6,016	5,990	5,450	256,053
Injection drug use	4,070	4,074	4,062	4,116	3,308	109,070
Male-to-male sexual contact and injection drug use	1,324	1,366	1,323	1,322	1,180	39,467
Heterosexual contact	1,389	1,528	1,513	1,634	1,548	24,268
Other ^b	195	166	167	160	113	9,843
Subtotal	12,933	13,202	13,080	13,222	11,599	438,701
Female adult or adolescent						
Injection drug use	1,892	1,907	1,977	1,989	1,744	41,178
Heterosexual contact	2,150	2,342	2,331	2,470	2,327	39,576
Other ^b	87	91	91	101	67	4,142
Subtotal	4,129	4,340	4,400	4,560	4,138	84,897
Child (<13 yrs at diagnosis)						
Perinatal	72	66	59	61	57	4,982
Other ^c	5	3	6	6	4	533
Subtotal	77	69	65	67	61	5,515
Region of residence						
Northeast	5,200	5,130	5,213	5,654	4,019	169,693
Midwest	1,622	1,646	1,623	1,199	1,234	50,333
South	7,078	7,386	7,361	7,839	7,192	181,690
West	2,567	2,683	2,585	2,428	2,540	108,183
U.S. dependencies, possessions, and associated nations	672	766	763	730	814	19,214
Total^d	17,139	17,611	17,544	17,849	15,798	529,113

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

^a Includes persons who died with AIDS, from the beginning of the epidemic through 2004.

^b Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes persons of unknown race or multiple races and persons of unknown sex. Cumulative total includes 836 persons of unknown race or multiple races. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.